

Options for Delivering Ecosystem-based
Marine Management

ODEMM



HUMBOLDT TIPPING

FRAMEWORK

03	Humboldt Tipping
04	Introduction
06	The ODEMM Framework
08	Methodology
10	Overview
12	I. Sectors
16	II. Pressures
20	III. Ecological Components

The Humboldt Tipping Project

The Humboldt Tipping project, with an interdisciplinary consortium of scientists from Germany and Peru, is focussed on understanding the risks posed by global change to the Humboldt Upwelling System and exploring potential adaptation options. In our second phase, we aim to integrate the insights gained from the first phase into a comprehensive analysis of future scenarios under the Shared Socio-economic Pathways (SSPs). This will help us to offer useful information for global climate change research and policy considerations.

To achieve this, we will engage closely with stakeholders: with those who know the Humboldt Upwelling System best.

Our central questions are:

1. What are the direct impacts on artisanal fishery communities?
2. What are the impacts on the industrial fleet?
3. How will the projected changes impact the local and global markets?
4. What kind of policy measures can ensure a sustainable use of the natural resources?

For more information visit:

<https://humboldt-tipping.org>



INTRODUCTION

Even for populations living far from the coast, the marine ecosystems play a critical role in every aspect of human life, by delivering multiple ecosystem services such as food production, climate regulation, participation in the nutrient cycle, and serving as a means of transport. In consequence, several marine ecosystems are among the most highly impacted on Earth by human activities. Additionally, the repercussions of climate change accelerate the degradation of marine ecosystems around the world.



© L. Kluger


This situation urged the creation of global legislation to attain sustainable management of the marine system and the preservation of ecosystem services.

Hence, the 2030 Agenda for Sustainable Development includes, among its 17 goals, a goal to “**Conserve and sustainably use the oceans, seas, and marine resources for sustainable development**”. Achieving this goal requires considering the whole suite of anthropogenic pressures affecting marine ecosystems. Yet the common perception is that fisheries are one of the biggest pressures on marine ecosystems in terms of magnitude and extension, although there are multiple other pressures that are less considered and should be tested to reduce the risk of pressures in marine ecosystems.

One way to test the anthropogenic pressures exerted by different sectors is to conduct **Integrated Ecosystem Assessments (IEA)**. IEAs propose ecosystem-based management based on a general view of coexisting pressures from identified sectors over ecosystem components. Hence, a key element of IEAs is the risk assessment of pressures on ecosystem components' susceptibility and resilience post-pressures. Among IEAs, there is a wide range of methodologies that can be classified as qualitative (based on expert references), quantitative assessments (based on monitoring and assessment of ecosystem indicators), and a mix of both. Qualitative and quantitative types of IEAs are complementary and after a qualitative IEA could be a quantitative being implemented for a selected number of indicators.

In the framework of the Humboldt Tipping project, we propose to use a qualitative approach to identify and assess the risk of multiple marine ecosystems that allows providing **Options for Delivering Ecosystem-based Marine Management (ODEMM)**.

The ODEMM framework



The ODEMM framework is a flexible tool that can be tailored to the characteristics of particular ecosystems and has integrated several requirements of legislative policy such as the Marine Strategy Framework Directive (MSFD; European Union, 2008), Common Fisheries Policy (CFP; European Union, 2013), and the Maritime Spatial Planning Directive (MSPD; European Union, 2014).

The main concept of ODEMM is the tracking of the pressures between multiple activities and the marine environment composed of ecological components, to provide a structure within which management options can be explored (Robinson et al., 2014). The ecological components are also linked with ecosystem services that are previously fixed using the reference of ODEMM typology (Böhnke-Henrichs et al., 2013).

Methodology

The methodology of ODEMM consists, as a first step, in the mapping of sectors-pressures-ecological components. This mapping will be carried out in 2 stages:

1. First, a pilot mapping of sectors-pressures-ecological components should be conducted among experts by means of an electronic survey.
2. During a second stage, the experts may gather in a panel assembled by scientists, advisors, policymakers, and members of the main sectors from national state agencies, scientific institutions, and companies with activities in marine ecosystems.

During the workshop, the pilot mapping of sectors-pressures-ecological components will be judged, and scores would be assigned for the spatial extent/overlap, frequency of occurrence, degree of impact, persistence, and resilience for each pressure-ecological component, based on predetermined categorical thresholds (Table 1).

The scores should be based in expertise and data and later supported by cross-checking using expert judgment and literature where gaps exist. Experts should judge and assign scores based on data. The experts will be classified into groups based on their knowledge of ecological components. The scores should be applied according to the current status (present facts) rather than potential risk in the future. A consensus for the mapping and scores is sought from each group of experts based on the best evidence or majority where consensus was not immediate. In the case of ecological components with multiple species, scoring should be based on assemblage and ecosystem functioning rather than focused on single species.

The qualitative scores should be later converted into numerical scores to calculate 'Proportional Connectance', and 'Impact Risk' (based on the 'overlap', 'frequency', and 'degree of impact' scores); and 'Recovery Lag' (based on 'resilience' and 'persistence' scores).

Proportional Connectance, Impact Risk, and Recovery Lag should serve to inform ecosystem management and evaluate the ecosystem components and ecological services paths at risk at present. Finally, the results should be related to the scenarios of climate change developed during the Humboldt tipping project and presented to the stakeholders in order to discuss options for marine ecosystem management in the future.

This booklet will give you an overview of the activities, sectors and ecological components that are important for this framework. If you are viewing it as a PDF, you can click on the text in the contents (page 02), overview (page 10) to get to the corresponding sections.

Table 1: Scores applied to each link between sectors- pressures- ecological components.

Spatial overlap	Frequency (annual)	Degree of impact	Resilience	Persistence
No overlap	Rare	Low	High	Low
Exogenous	Occasional	Chronic	Moderate	Moderate
Site	Common	Acute	Low	High
Local	Persistent		None	Continuous
Widespread patchy				
Widespread even				

Overview

I. Sectors

Aquaculture
Artisanal fisheries
Industrial fisheries
Maritime transport

Renewable energy
Oil and gas
Coastal infrastructure
Coastal industry

Agriculture
Tourism and recreation
Domestic sewage

II. Pressures

Physical loss

Sealing

Physical disturbance

Noise
Litter

Physical damage

Abrasion
Smothering and turbidity
Extraction of non-biological resources

Hydrological process interference

Thermal regime
Wave exposure and current changes

Chemical composition interference

Salinity regime
pH changes
Contaminating compounds
Organic matter

Biological disturbance

Invasive species
Barriers
Species extraction
Bycatch
Incidental loss

III. Ecological Components

Shelf

Peruvian anchovy
Coastal pelagic fish
Shelf pelagic
Coastal demersal fish
Shelf sediment
Shelf rock
Scallops
Demersal elasmobranchia

Littoral

Littoral sediment
Littoral rock
Estuaries
Seabirds and mammals

Oceanic

Giant squid
Oceanic pelagic fish
Pelagic elasmobranch
Oceanic pelagic
Deep sea bottom

Disclaimer

Thanks for your participation in the ODEMM survey. This survey is part of the research activities of the Humboldt tipping project. Here we detailed the conditions of data usage:

1. The data collected will be used to better understand the impacts on marine ecosystems and to formulate management fisheries recommendations.
2. The data provided will be treated anonymously and will not be used for any other purpose outside of the project.
3. Upon completion of the study, a summary of the results will be communicated to you.
4. No monetary compensation is offered for your participation in this survey.
5. If at any time you decide not to continue, you can close the app.
6. This booklet was created by Lena Hindenberg and Giannina Passuni in the frame of Humboldt Tipping, and is protected by copyright.
7. For any question and detailed information about this study, please contact us at: giannina.passuni.saldana@uni-hamburg.de



Giannina Passuni,
University of Hamburg



I. Sectors

Aquaculture



The main cultivation resource is the calico scallop, which is mainly found in the bays of Sechura, Samanco, and Pisco. Scallops are raised in floating cages that are constantly monitored by boats and diving. Other important crops include macroalgae and shrimp farming in mangroves. The aquaculture sector is still in development and had potential to expand production and diversify the species cultivated. Obstacles to develop the aquaculture are infrastructure, regulation, and staff training.

Artisanal fisheries



Artisanal fishing in Peru involves small-scale operations using vessels up to 15 m long and 32 m³ hold capacity. It operates within 5 miles from the coast, focusing on meeting local fresh and frozen fish demand.

The sector grew 300% from 1996 to 2015, but now faces challenges like oversized fleets, overfishing, and regulatory issues, leading to conflicts with industrial fisheries. Effective management policies are crucial for the long-term viability of the sector.

Industrial fisheries



Industrial fishing is the second largest sector contributing to the Peruvian GDP. The only species targeted by this sector is Peruvian anchovy for the production of fish oil and meal. Around 6 million tons of fish are caught annually, yielding 1.5 million tons of fishmeal and 150,000 tons of oil, mainly exported to China (78%), Ecuador (6.2%), and Germany (5.4%). The anchovy fishing is carried out by around 800 metal and wood vessels operating with purse seine nets. It's dependency on a single species poses challenges for sustainability.

Maritime transport



Maritime transport is vital for Peru's economic growth, serving as a key link for imports & exports, mainly from the ports Paita, Salaverry, Chimbote, Callao, Pisco, Matarani and Ilo. With strategic positioning between the Pacific with connections to the Atlantic, Peru handles approximately 97 million tons of cargo annually, with 91% of exports shipped by sea. While trade volumes are rising, there is a need for better port management, fleet modernization, and infrastructure upgrades. Balancing development with sustainability requires addressing environmental impacts. Passenger transport, notably large cruise ships is still in development.

Renewable energy



Renewable energy, like wind and tidal power, holds promise for sustainable energy in Peru. Despite potential coastal areas, due to limited investment and technological development, only one pilot wind energy program exists in San Juan de Marcona. Proper marine space planning is crucial to prevent conflicts with other sectors, and a robust regulatory framework is needed to address environmental risks associated with large-scale projects.

Oil and gas



Oil and gas industry are widely developed along the coast and the continental shelf, contributing to 24% of the national fuel production (20 million barrels annually; World Bank). Activities in the maritime scope of the northern and central coast include seismic exploration, exploitation, and refining. The main refineries are Talara, La Pampilla, and Conchan. Activities of this sector can pose environmental risks, as seen in the recent oil spill off La Pampilla.

Coastal infrastructure



This sector includes all forms of infrastructure on the coastal edge such as ports, resorts, breakwaters, and beach nourishment. It also includes coastal edge modification, for example, land reclamation through the drying of wetlands for use in agriculture or housing. Peru had 45 maritime ports and the most important are Paita, Salaverry, Chimbote, Callao, Pisco, Matarani, and Ilo. Several of the biggest cities are close to the coast and includes coastal resorts as Punta Sal (Tumbes) and Huanchaco (La Libertad), which have undergone refurbishment or expansion due to coastal erosion and increased tourism. The growth of this sector has faced challenges like sustainable management of coastal resources and adaptation to climate change.

Coastal industry



Activities included here are within a 5km radius of the coastal edge and should interact with the marine environment, either through material extraction or the discharge of products into the sea. Examples of important industries on the Peruvian coast include fish processing companies, canneries (Tumbes and Piura), fish meal producers (Chimbote & Pisco), desalination plants (PROVISUR), mining (Shogun Iron Mine in Ica), and poultry farms (90% of national poultry production occurs on the coast).

Agriculture



Traditional crops on the Peruvian coast include cotton, sugarcane, and rice. However, since 2000, agriculture had a large development through the diversification of crops including export fruits such as blueberries, avocados, and asparagus. Agriculture can interact with the marine environment through runoff and drainage. Although the water deficit in mostly sandy terrain decrease the interaction with marine areas, except during torrential rains due to El Nino events. Rains produces landslides known as "huaycos", which increase runoff of sediments and organic matter, including chemical compounds from fertilizers and pesticides to the sea.

Tourism and recreation



Constituting the third largest sector contributing the most to the Peruvian GDP. In the marine environment, tourism focuses on beaches and resorts in the summer. Marine reserves, such as the Paracas National Reserve attract visitors through their rich marine and terrestrial biodiversity. Tourism activities include the use of recreational boats, surfing, recreational and sports fishing, water parks, horseback riding or walking along beaches, and food vending. Tourism exerts significant pressures on the marine environment, including water pollution, disturbances of habitats, vessel collisions with marine species, and overexploitation of resources.

Domestic sewage




In Peru, there is a deficit in wastewater treatment. According to the entity overseeing the operation and maintenance of sewage systems (Sunass), 77% of wastewater is treated in plants located in major cities. From them, 11% of sewage after treatment in plants is discharged into the sea. Although the level of treatment in most of the plants only includes the removal of macro-residues.



II. Pressures

Physical loss

Sealing



The physical loss of soil due to sealing by construction or permanent substrate change. It becomes permanent when structures are built for longer than 50 years. Soil includes substrate from the beach, intertidal & subtidal bottom, and seabed.

Sources: port construction, beach urbanization, installation of offshore wind parks

Physical disturbance

Noise



The introduction of sounds into the aquatic environment can interfere with the communication and navigation of marine species. Birds, marine mammals, and fish depend on sound for feeding, orientation, and communication. Noise pollution can disrupt their behavioral patterns.

Sources: boat engines, acoustic surveys, underwater construction, and military activities


Litter



Litter generated by domestic activities, fishing, agriculture, and coastal industry that enter the marine environment. The litter considered includes microplastics, macroplastics, metals, glass, rubber, wood, and fabric. Only the direct effects on marine organisms are considered, excluding the effects on the food chain.

Physical damage

Abrasion



The wear or removal of the seabed due to human activities such as, anchoring, trawling, and removal using air compressors. The substrate includes the beach, the intertidal and subtidal zones, and the seabed.

Sources: dredging for the construction of ports, navigation channels and anchoring of vessels


Smothering and turbidity



The increase in sediments in the water column can hinder, among other factors, the penetration of sunlight, negatively affecting photosynthesis. Additionally, turbidity and sedimentation on the seafloor can impede oxygen circulation and clog gills and filtering apparatuses, causing suffocation and death of organisms like mollusks, crustaceans, and fish.

Sources: construction of coastal infrastructure, and runoff from agriculture and coastal industry

Extraction of non-biological resources



Involves the removal or collection of materials from water column and seafloor for industrial use. This includes mining, guano extraction, and the extraction of sand, gravel, and seawater. Resource extraction can alter sedimentation patterns, water quality, and have adverse effects on coastal stability, biodiversity, and habitat conservation.

Sources: seawater extraction for industrial plant cooling, desalination, sand and gravel extraction for construction, and concrete production

Hydrological process interference

Thermal regime

Variations in water temperature, both in average and range, due to drainage discharge. This pressure does not include variations due to natural oceanographic phenomena or climate change. Increased water temperature can cause thermal stress in ectothermic organisms, resulting in reduced growth and survival.

Sources: industrial, urban, and agricultural wastewater, cooling water from power plants

Wave exposure and current changes

Modifications in water dynamics, such as currents and tides, caused by the presence of coastal infrastructure. These changes can alter the water flow, nutrient distribution, and sedimentation for a given area. This can especially affect the organisms that cannot actively move, like plankton, macroalgae, and fish larvae. They depend on currents for their dispersion.




Sources: docks, dams, dikes, wind turbines, and other artificial structures

Chemical composition interference

pH changes

Alterations in the acidity or alkalinity level of seawater and/or substrate. This excludes oceanographic changes, natural river flow, and changes caused by climate change.

Sources: industrial and domestic wastewater, and changes in substrate structure



Salinity regime

Variations in the concentration of dissolved salts in seawater, both in average and range, due to human activities. This pressure does not include variations due to natural oceanographic phenomena or climate change. Stenohaline organisms can be affected, leading to stress and changes in distribution.

Sources: domestic and industrial wastewater, and alteration in the mixing of fresh and salt-water due to coastal infrastructure

Contaminating compounds




The incorporation of man-made substances, like pesticides, pharmaceuticals, heavy metals, and hydrocarbon compounds in the water column or marine sediment. For example, pesticides in agricultural activities can be carried by runoff water, pharmaceuticals by urban wastewater, and hydrocarbons by marine transport accidents. Heavy metals, like mercury, lead, and cadmium from mining can also accumulate in marine organisms.

Sources: agriculture, industry, fishing, oil and gas industry, and domestic use

Organic matter

The enrichment of the water column and marine sediment with organic materials, which can lead to water deoxygenation. For example, the discharge of organic waste, such as fish remains, and unconsumed food can decompose and consume oxygen. This can negatively affect marine life.

Sources: agricultural runoff, urban and industrial drainage, aquaculture effluents



Biological disturbance

Invasive species

The introduction of non-native organisms as for example, the dinoflagellates *Alexandrium ostenfeldii*, introduced through ballast water in the Peruvian coast. Dinoflagellates can trigger harmful algal blooms, which have adverse effects on the health of ecosystems, marine life, and human activities.

Sources: ballast water exchange of ships, aquaculture, and wastewater

Barriers



Limitation in the movement of marine fauna along a key migration route or inside its distribution habitat. The formation of barriers due to human activities can impact life history processes such as reproduction, feeding, or seasonal migration. This considers species that actively move, such as fish, birds, and marine mammals.

Sources: presence of coastal infrastructure, such as piers, dikes, and turbines

Species extraction

The collection or capture of biological resources from the beach, water column, or seabed. These activities may involve extracting parts of an individual, capturing the entire individual, or even entire populations. The extraction of biological resources has secondary effects on the population dynamics of species. When it reaches the point of overexploitation, it can alter the population viability of one or multiple species.

Sources: the extraction of marine biological resources such as fish, mollusks, and crustaceans





Bycatch

Unintentional capture of species during fishing activities, usually of fish and invertebrates. Incidental capture can be associated with the low specificity of fishing nets. The pressure from incidental capture often leads to the discarding of these species or their subsequent use for consumption or commercialization. Bycatch can include the capture of endangered or protected species such as sharks and dolphins which are later commercialized.

Incidental loss

Death or injury of marine fauna due to human activities. This would primarily affect top predators and charismatic species such as birds, whales, dolphins, and sea turtles. For example, entanglement in fishing nets and fishing gears can cause serious injury and/or subsequent mortality. Additionally, collisions with vessels pose a significant threat to the conservation of these species, as they can result in severe injury or even death following an accident.





III. Ecological Components

Shelf

Peruvian anchovy

Peruvian anchovy is a pelagic fish that inhabits the cold waters of the Humboldt Current and is key in the food chain for multiple fish, birds, and mammals. It has great commercial importance, with around 6 million tons of annual landings, making it the largest single-species fishery. Despite having the highest biomass in the pelagic system, it is vulnerable to climatic variability and overfishing. Locations: north-central stock off the Peruvian coast & southern stock, shared with Chile.

Coastal pelagic fish

Includes other bony fish present in the water column within the continental shelf (<200 nautical miles), with the exception of anchovy.

Among the commercial species of pelagic fish are mackerel (*Scomber japonicus*), jack mackerel (*Trachurus murphyi*), and bonito (*Sarda chilensis*). The capture of these species is mainly carried out by artisanal fishing and destined for direct consumption.

Shelf pelagic

Includes pelagic elements, both abiotic (seawater, currents, nutrients) and biotic, distributed within the continental shelf (<200 nautical miles). The biotic elements include coastal plankton, composed of phytoplankton (20 μm - 200 μm), zooplankton (200 μm - 2 μm), and jellyfish (2 cm - >1 m). Pelagic fish described in an individual category are excluded.

Coastal demersal fish

Bony (demersal) fish that inhabit and are associated with the seafloor for food or shelter, distributed mainly in coastal waters (<200 m depth). They are found either on soft substrates (sand or mud) or rocky substrates and caught for direct consumption by artisanal fishing. Among commercially important species are: kelp bass (*Paralabrax humeralis*), and grouper (*Mycteroperca xenarcha*).

Shelf sediment

Characterized by sandy muddy substrate extending from the subtidal zone to the edge of the continental shelf (<200 nautical miles). It is defined by oxygen levels that allow the distribution of organisms sensitive to hypoxia and where light penetration allows the growth of more plankton than at the bottom. It hosts, for example, polychaetes, mollusks, bivalves, and crustaceans, like the clam (*Leukoma thaca*), razor clam (*Ensis macha*), and the moon snail (*Sinum cymba*).

Shelf rock

Characterized by the presence of benthic rocky substrates, including natural outcrops, artificial reefs, or artificial structures. It extends from the subtidal zone to the continental shelves, covering a range from a few meters to 200 m. It hosts a variety of species like polychaetes, mollusks, bivalves, and crustaceans. Some commercially important species are: black mussels (*Aulacomya atra*), hairy crab (*Romaleon setosum*), and green sea urchin (*Loxechinus albus*).

Scallops


A commercially important bivalve mollusk. It is mainly found on the central and southern coast, especially in the bays of Paracas, Pisco, and Independencia. Its ideal habitat is cold, nutrient-rich waters with sandy or muddy bottoms between 10-30 m. The scallop fishery depends mainly on cultivation by catching seeds from natural beds and larvae. It has high international demand. In 2021, scallop export value reached 122 million dollars.

Demersal elasmobranchia

Cartilaginous fish associated with the bottom for food and shelter. They are mostly distributed in coastal waters (>200 m depth). They are targeted for local consumption, however, many species are threatened by overfishing. The most characteristic species are common dogfish (*Mustelus whitneyi*), angel shark (*Squatina californica*), and common guitarfish (*Pseudobatos planiceps*).

Littoral

Littoral sediment



It is a coastal ecosystem composed by sandy substrate, covering the supratidal and intratidal zones. Some examples of organisms found here are polychaetes, crustaceans, and mollusks. Some characteristic crustaceans are


Emerita analoga and *Neotrypaea californiensis* often used as bait in artisanal & recreational fishing. Also the mollusk Peruvian bean clam (*Donax peruvianus*) and concha macha (*Mesodesma donacium*) are commercially important.



Seabirds and mammals


Includes birds inhabiting beaches, estuaries, and islands, and feeding on marine resources. It also includes marine mammals that reside on land and feed in coastal waters, and marine pelagic mammals residents. Peru has around 60 species of birds, including 3 which are known for their guano production: Guanay cormoran (*Leucocarbo bougainvillii*), the Peruvian booby (*Sula variegata*) and the Peruvian pelican (*Pelecanus thagus*). Some marine mammals are the south american sea lion (*Otaria flavescens*), the south american fur seal (*Arctocephalus australis*), and the bottlenose dolphin (*Tursiops truncatus*).

Littoral rock



Characterized by the presence of rocky substrates emerging from the sea, either permanently or during low tide, ranging from pebbles to larger rocks, extending from the supratidal to the rocky intertidal zone. They provide a unique habitat to species of macroalgae (for example, *Ulva lactuca*), bivalves (for example, *Perumytilus purpuratus*), and commercially important molluscs like limpets (*Fissurella spp.*) and goose barnacles (*Pollicipes elegans*).

Example: Ancon Bay



Estuaries

Coastal ecosystems placed in the transition between brackish water and the sea. Mangroves consist of salt-tolerant vegetations. They are important for coastal erosion control, carbon storage, and particle filtration. They provide a habitat for species like the red mangrove crab (*Ucides cordatus*) and black clam (*Anadara chemnitzii*).


Estuaries are semi-closed coastal ecosystems where river water mixes with seawater, influences by tides & currents. They serve as fish nurseries, nesting sites for birds, and resting places of species like the green turtle (*Chelonia mydas agassizii*).

Wetlands include swamps and ecosystems permanently flooded by brackish water from rain, rivers, or groundwater. They are carbon and organic matter sinks, filter particles, and regulate tides. They are fish breeding grounds and habitats to species like waterfowl.

Examples: regions of Tumbes & Piura (mangroves), Virrila estuary & Cira River estuary (estuaries), and Pantanos de Villa & Albufera de Medio Mundo in Lima (wetlands)


Oceanic

Giant squid




Dosidicus gigas is a cephalopod that can reach up to 1.8 meters in length and weigh up to 50 kg. It is distributed in the temperate and subtropical waters of the eastern Pacific Ocean, from the United States to Chile. In Peru, the giant squid is one of the fishing species with an annual catch of around 500,000 tons. Landings are used both for local consumption and for export.

Oceanic pelagic fish




This category includes pelagic and migratory bony fish that commonly inhabit oceanic waters and only occasionally approach the Peruvian coast. Amongst the most important are dolphin fish, tuna fish, swordfish, and flying fish. The dolphin fish is the 3rd most important target for artisanal fishery using surface longlines and curtain nets. Despite a Fisheries Management Regulation for sustainable fishing of dolphin fish only took effect in January 2024. Tuna fishing is regulated by the Inter-American Tropical Tuna Commission (IATTC) resolution.

Pelagic elasmobranchia




Includes cartilaginous fish that are distributed in the water column. While typically distributed in oceanic waters they can also enter coastal waters. Among the species most important are thresher shark, manta ray, shortfin mako shark, and blue shark. These top predators are crucial in the food chain and commercially important for consumption and fin extraction. Overfishing threatens many species, prompting a finning ban. CITES has advised Peru to strengthen shark and ray trade regulations, as Peru is the largest shark fin exporters in Latin-America.

Oceanic pelagic



Includes various pelagic elements, both abiotic (seawater, currents, nutrients) and biotic components. These include phytoplankton (20 μm - 200 μm), zooplankton (200 μm - 2 μm), and jellyfish (2 cm - >1 m). They are distributed beyond the continental shelf (>200 nautical miles), it also encompasses migratory megafauna like humpback and blue whales, though fish are classified differently.

Deep sea bottom



Extends from the continental shelf edge to the seabed, including abiotic components like sediments, rocks, and minerals (e.g. manganese nodules and hydrothermal deposits) and biotic elements like abyssal fish. These include blackbelly rosefish, lantern fish and genadier. These resources are not currently exploited off Peru's coast.



Kiel University
Christian-Albrechts-Universität zu Kiel



Universität Hamburg
DER FORSCHUNG | DER LEHRE | DER BILDUNG



University
of Bremen



LEIBNIZ CENTRE
for Tropical Marine Research



HUMBOLDT
TIPPING



GEOMAR



IMARPE
INSTITUTO DEL MAR DEL PERÚ



PUCP



Federal Ministry
of Education
and Research

FONA

Forschung für Nachhaltigkeit